

Lampiran 4. Hasil Uji Statistik Eritrosit

A. Uji Normalitas

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
eritrosit	,076	48	,200*	,977	48	,467

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

B. Uji Homogenitas

Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
eritrosit	Based on Mean	,978	5	42	,442
	Based on Median	,665	5	42	,652
	Based on Median and with adjusted df	,665	5	33,005	,653
	Based on trimmed mean	,932	5	42	,470

C. Uji Two-Way Anova

Tests of Between-Subjects Effects

Dependent Variable: eritrosit

Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Squared
Corrected Model	12,960 ^a	11	1,178	2,154	,041	,397
Intercept	2256,255	1	2256,255	4125,257	,000	,991
kelompok	4,337	5	,867	1,586	,189	,181
perlakuan	7,061	1	7,061	12,910	,001	,264
kelompok * perlakuan	1,562	5	,312	,571	,721	,074
Error	19,690	36	,547			
Total	2288,905	48				
Corrected Total	32,650	47				

a. R Squared = ,397 (Adjusted R Squared = ,213)

Lampiran 5. Hasil Uji Statistik Hemoglobin

A. Uji Normalitas

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
hemoglobin	,131	48	,039	,972	48	,315

a. Lilliefors Significance Correction

B. Uji Homogenitas

Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
hemoglobin	Based on Mean	,922	5	42	,476
	Based on Median	,899	5	42	,491
	Based on Median and with adjusted df	,899	5	34,370	,493
	Based on trimmed mean	,909	5	42	,485

C. Uji Two-Way Anova

Tests of Between-Subjects Effects

Dependent Variable: hemoglobin

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	44,767 ^a	11	4,070	1,814	,088
Intercept	8279,253	1	8279,253	3690,603	,000
kelompok	14,547	5	2,909	1,297	,287
perlakuan	21,333	1	21,333	9,510	,004
kelompok * perlakuan	8,887	5	1,777	,792	,562
Error	80,760	36	2,243		
Total	8404,780	48			
Corrected Total	125,527	47			

a. R Squared = ,357 (Adjusted R Squared = ,160)

Lampiran 6. Hasil Uji Statistik Hematokrit

A. Uji Normalitas

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
hematokrit	,083	48	,200*	,977	48	,475

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

B. Uji Homogenitas

Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
hematokrit	Based on Mean	,923	5	42	,476
	Based on Median	,884	5	42	,500
	Based on Median and with adjusted df	,884	5	30,933	,504
	Based on trimmed mean	,931	5	42	,471

C. Uji Two-Way Anova

Tests of Between-Subjects Effects

Dependent Variable: hematokrit

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	358,057 ^a	11	32,551	1,931	,068
Intercept	56842,567	1	56842,567	3372,696	,000
kelompok	91,292	5	18,258	1,083	,386
perlakuan	201,720	1	201,720	11,969	,001
kelompok * perlakuan	65,045	5	13,009	,772	,576
Error	606,735	36	16,854		
Total	57807,360	48			
Corrected Total	964,792	47			

a. R Squared = ,371 (Adjusted R Squared = ,179)

