

UJI STATISTIKA SPSS

1. Uji normalitas shaphiro-wilk

Tests of Normality

		Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	kelompok	Statistic	df	Sig.	Statistic	df	Sig.
kadar	jarak	0,125	16	0,200 [*]	0,925	16	0,206
	perokok	0,193	16	0,115	0,936	16	0,308

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

2. Uji homogenitas Lavene

Levene's Test of Equality of Error Variances^{a,b}

		Levene Statistic	df1	df2	Sig.
kadar	Based on Mean	1,812	1	30	0,188
	Based on Median	1,985	1	30	0,169
	Based on Median and with adjusted df	1,985	1	29,999	0,169
	Based on trimmed mean	1,762	1	30	0,194

Tests the null hypothesis that the error variance of the dependent variable is equal across groups.

a. Dependent variable: kadar

b. Design: Intercept + kelompok

3. Uji Pengaruh Two way anova

Tests of Between-Subjects Effects

Dependent Variable: kadar

Source	Type III Sum of Squares	df	Mean Square	F	Sig.
Corrected Model	7,498E-7 ^a	1	7,498E-7	0,128	0,723
Intercept	0,007	1	0,007	1279,660	0,000
kelompok	7,498E-7	1	7,498E-7	0,128	0,723
Error	0,000	30	5,856E-6		
Total	0,008	32			
Corrected Total	0,000	31			

a. R Squared = ,004 (Adjusted R Squared = -,029)