

## Lampiran 2. Hasil Uji Statistik

- Uji Normalitas Shapiro-Wilk

Tests of Normality						
	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
TROMBOSIT	.134	30	.182	.942	30	.105
HEMATOKRIT	.132	30	.192	.942	30	.103
ALT	.146	30	.100	.898	30	.008
BILIRUBINT	.401	30	.000	.397	30	.000

a. Lilliefors Significance Correction

KODE	LOG10 ALT	LOG10 BILT
1.	1.79	-.15
2.	2.05	-.30
3.	1.69	-.52
4.	1.80	-.30
5.	1.64	-.40
6.	2.14	-.40
7.	1.83	-.22
8.	1.89	-.40
9.	1.53	.43
10.	1.36	-.52
11.	1.65	-.40
12.	2.02	-.52
13.	1.00	-.52
14.	1.23	-.40
15.	1.74	-.52

KODE	LOG10 ALT	LOG10 BILT
16.	1.43	-.10
17.	1.49	-.40
18.	1.20	-.40
19.	1.68	-.15
20.	1.96	-.15
21.	1.08	.83
22.	1.30	-.40
23.	1.92	-.10
24.	1.64	.04
25.	1.04	-.22
26.	1.61	-.22
27.	1.23	-.22
28.	1.26	-.10
29.	1.53	-.70
30.	1.40	-.15

- Uji Normalitas Shapiro-Wilk setelah LOG10

Tests of Normality						
	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
TROMBOSIT	.134	30	.182	.942	30	.105
HEMATOKRIT	.132	30	.192	.942	30	.103
LOG_ALT	.090	30	.200 <sup>*</sup>	.974	30	.664
LOG_BILIRUBINT	.200	30	.003	.816	30	.000

\*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

- Uji Korelasi Pearson Trombosit, Hematokrit, ALT

Correlations					
		TROMBOSIT	HEMATOKRIT	LOG_ALT	BILIRUBINT
TROMBOSIT	Pearson Correlation	1	-.450 <sup>*</sup>	-.549 <sup>**</sup>	.218
	Sig. (2-tailed)		.013	.002	.247
	N	30	30	30	30
HEMATOKRIT	Pearson Correlation	-.450 <sup>*</sup>	1	.230	.136
	Sig. (2-tailed)	.013		.222	.472
	N	30	30	30	30
LOG_ALT	Pearson Correlation	-.549 <sup>**</sup>	.230	1	-.280
	Sig. (2-tailed)	.002	.222		.134
	N	30	30	30	30
BILIRUBINT	Pearson Correlation	.218	.136	-.280	1
	Sig. (2-tailed)	.247	.472	.134	
	N	30	30	30	30

\*. Correlation is significant at the 0.05 level (2-tailed).

\*\*. Correlation is significant at the 0.01 level (2-tailed).

- Uji Korelasi Spearman Bilirubin Total

**Correlations**

			TROMBOSIT	HEMATOKRIT	LOG_ALT	BILIRUBINT
Spearman's rho	TROMBOSIT	Correlation Coefficient	1.000	-.378*	-.502**	.112
		Sig. (2-tailed)	.	.039	.005	.554
		N	30	30	30	30
	HEMATOKRIT	Correlation Coefficient	-.378*	1.000	.245	.181
		Sig. (2-tailed)	.039	.	.193	.338
		N	30	30	30	30
	LOG_ALT	Correlation Coefficient	-.502**	.245	1.000	-.054
		Sig. (2-tailed)	.005	.193	.	.777
		N	30	30	30	30
	BILIRUBINT	Correlation Coefficient	.112	.181	-.054	1.000
		Sig. (2-tailed)	.554	.338	.777	.
		N	30	30	30	30

\*. Correlation is significant at the 0.05 level (2-tailed).

\*\*. Correlation is significant at the 0.01 level (2-tailed).