

## Hasil Analisis SPSS

### Uji Normalitas

#### Tests of Normality

	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
HD	.057	204	.099	.992	204	.370
IF	.073	204	.011	.989	204	.112
PD	.076	204	.007	.989	204	.130

a. Lilliefors Significance Correction

### Uji Linieritas

#### ANOVA<sup>a</sup>

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	1598.863	2	799.432	183.549	.000 <sup>b</sup>
	Residual	875.439	201	4.355		
	Total	2474.302	203			

a. Dependent Variable: PD

b. Predictors: (Constant), IF, HD

### Uji Korelasi Pearson

#### Correlations

		HD	IF	PD
HD	Pearson Correlation	1	.766**	.800**
	Sig. (2-tailed)		.000	.000
	N	204	204	204
IF	Pearson Correlation	.766**	1	.662**
	Sig. (2-tailed)	.000		.000
	N	204	204	204
PD	Pearson Correlation	.800**	.662**	1
	Sig. (2-tailed)	.000	.000	
	N	204	204	204

\*\*. Correlation is significant at the 0.01 level (2-tailed).

### Uji R Square

#### Model Summary<sup>b</sup>

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.804 <sup>a</sup>	.646	.643	2.087

a. Predictors: (Constant), IF, HD

b. Dependent Variable: PD