

TABEL DISTRIBUSI FREKUENSI

Pretest

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	50	2	11.8	11.8	11.8
	65	2	11.8	11.8	23.5
	70	6	35.3	35.3	58.8
	75	5	29.4	29.4	88.2
	80	2	11.8	11.8	100.0
	Total	17	100.0	100.0	

Posttest

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	70	1	5.9	5.9	5.9
	75	1	5.9	5.9	11.8
	80	3	17.6	17.6	29.4
	85	4	23.5	23.5	52.9
	90	3	17.6	17.6	70.6
	95	5	29.4	29.4	100.0
	Total	17	100.0	100.0	

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
Pretest	17	50	80	69.71	8.564
Posttest	17	70	95	86.47	7.658
Valid N (listwise)	17				

HASIL UJI NORMALITAS DATA

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Pretest	.278	17	.001	.816	17	.003
Posttest	.161	17	.200 [*]	.905	17	.082

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

HASIL UJI T SAMPEL BERPASANGAN (*PAIRED T TEST*)

Paired Samples Statistics

	Mean	N	Std. Deviation	Std. Error Mean
Pair 1 Pretest	69.71	17	8.564	2.077
Posttest	86.47	17	7.658	1.857

Paired Samples Correlations

	N	Correlation	Sig.
Pair 1 Pretest & Posttest	17	.484	.049

Paired Samples Test

		Paired Differences					t	df	Sig. (2-tailed)
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	Pretest - Posttest	-16.765	8.280	2.008	-21.022	-12.507	-8.348	16	.000