

## 2. Hasil Uji Data Statistik SPSS Versi 27

### a. Uji Normalitas dan Homogenitas

Tests of Normality							
Kelompok Pasien		Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Eritrosit	DM	,159	30	,052	,938	30	,079
	DM CKD	,124	30	,200 <sup>*</sup>	,967	30	,458
HBA1c	DM	,151	30	,080	,932	30	,056
	DM CKD	,117	30	,200 <sup>*</sup>	,931	30	,051
HB	DM	,129	30	,200 <sup>*</sup>	,956	30	,250
	DM CKD	,131	30	,200	,974	30	,657

\*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Test of Homogeneity of Variance					
		Levene Statistic	df1	df2	Sig.
Eritrosit	Based on Mean	,601	1	58	,441
	Based on Median	,560	1	58	,457
	Based on Median and with adjusted df	,560	1	48,212	,458
	Based on trimmed mean	,613	1	58	,437
HBA1c	Based on Mean	,066	1	58	,799
	Based on Median	,135	1	58	,715
	Based on Median and with adjusted df	,135	1	53,409	,715
	Based on trimmed mean	,067	1	58	,797
HB	Based on Mean	2,916	1	58	,093
	Based on Median	2,910	1	58	,093
	Based on Median and with adjusted df	2,910	1	52,200	,094
	Based on trimmed mean	2,932	1	58	,092

### b. Uji Independent Sample Test

Independent Samples Test									
		Levene's Test for Equality of Variances						t-test for Equality of Means	
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference
Eritrosit	Equal variances assumed	,601	,441	6,611	58	<,001	1,06833	,16160	,74486 1,39181
	Equal variances not assumed			6,611	53,614	<,001	1,06833	,16160	,74430 1,39237